


Critical-Care Observation Tool (CPOT)

Indicator	Score	Description	
Facial expression  Caroline Arbour, RN, B.Sc., PhD(student) School of Nursing, McGill University	Relaxed, neutral	0	No muscle tension observed
	Tense	1	Presence of frowning, brow lowering, orbit tightening and levator contraction or any other change (e.g. opening eyes or tearing during nociceptive procedures)
	Grimacing	2	All previous facial movements plus eyelid tightly closed (the patient may present with mouth open or biting the endotracheal tube)
Body movements	Absence of movements or normal position	0	Does not move at all (doesn't necessarily mean absence of pain) or normal position (movements not aimed toward the pain site or not made for the purpose of protection)
	Protection	1	Slow, cautious movements, touching or rubbing the pain site, seeking attention through movements
	Restlessness/Agitation	2	Pulling tube, attempting to sit up, moving limbs/thrashing, not following commands, striking at staff, trying to climb out of bed
Compliance with the ventilator (intubated patients) OR Vocalization (extubated patients)	Tolerating ventilator or movement	0	Alarms not activated, easy ventilation
	Coughing but tolerating	1	Coughing, alarms may be activated but stop spontaneously
	Fighting ventilator	2	Asynchrony: blocking ventilation, alarms frequently activated
	Talking in normal tone or no sound	0	Talking in normal tone or no sound
	Sighing, moaning	1	Sighing, moaning
	Crying out, sobbing	2	Crying out, sobbing
Muscle tension Evaluation by passive flexion and extension of upper limbs when patient	Relaxed	0	No resistance to passive movements
	Tense, rigid	1	Resistance to passive movements