Pediatric Pain and Fever Liquid Dosing Guide*

Acetaminophen (Tylenol®) **Dosing Table** (give every 4-6 hours as directed)									
Child's Age	0-3 mo*	4-11 mo	12-23 mo	2-3 yr	4-5 yr	6-8 yr	9-10 yr	11-12 yr	12+ yr
Weight (pounds)	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	96+ lbs
Weight (kilograms)	3-5 kg	6-7 kg	8-10 kg	11-15 kg	16-21 kg	22-26 kg	27-32 kg	33-43 kg	44+ kg
Liquid 160mg/5mL (mL)	1.25 mL	2.5 mL	3.75 mL	5 mL	7.5 mL	10 mL	12.5 mL	15 mL	20 mL

* **Speak to your doctor before giving Acetaminophen to children under 2 months old**

Ibuprofen (Advil®/Motrin®) **Dosing Table** (give every 6-8 hours as directed)									
Child's Age	0-6 mo	6-11 mo	12-23 mo	2-3 yr	4-5 yr	6-8 yr	9-10 yr	11-12 yr	12+ yr
Weight (pounds)	0-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	96+ lbs
Weight (kilograms)	0-5 kg	6-7 kg	8-10 kg	11-15 kg	16-21 kg	22-26 kg	27-32 kg	33-43 kg	44+ kg
Drops 50mg/1.25mL (mL)	--	1.25 mL	1.875 mL	2.5 mL	3.75 mL	5 mL	--	--	--
Liquid 100mg/5mL (mL)	--	2.5 mL	4 mL	5 mL	7.5 mL	10 mL	12.5 mL	15 mL	20 mL

* Ideal dosing is based on **weight**, not age. Use a dosing cup or syringe if possible. 1 teaspoon = 5 mL

mo = months of age; **yr** = years of age

pami.emergency.med.jax.ufl.edu

UF Department of Emergency Medicine
College of Medicine – Jacksonville
UNIVERSITY of FLORIDA

PAMI Pain Assessment and Management Initiative