# About OTC Topical Pain Medications

- Medications available for purchase at local pharmacies and stores without a prescription are called over-the-counter (OTC) medications
- OTC topical medications are applied to the skin and can help pain by numbing or reducing inflammation in the joints
- OTC topical pain relievers are available as creams, gels, lotions, sprays, and patches
- Common OTC topical medications include:
  - lidocaine
  - nonsteroidal anti-inflammatory drugs (NSAIDs)
  - counterirritants (capsaicin, menthol, camphor)
- Ask your local pharmacist if the cost of these medications is covered by your insurance plan

#### Warnings

- Read and follow all directions and warnings on packaging
- Review all active ingredients in the product Many products have multiple active ingredients
- It is important to use these medications only as directed. Stop using the product if any side effects occur and seek medical attention
- Keep out of reach of children and pets.
- See package for proper disposal



## **Contact us**

Phone: 904 - 244 - 4986 Email: pami@jax.ufl.edu

Website: pami.emergency.med.jax.ufl.edu

Search us using: @ufpami





# Scan the QR code to visit our website:





# OVER-THE-COUNTER TOPICAL PAIN MEDICATIONS

A guide to understanding topical pain medications



Always tell your healthcare provider about all medications you are taking, including OTC options, herbals and supplements.

## **Lidocaine 4% Patches**

Numerous brand names

### Diclofenac 1% Gel

Known as an NSAID

Example brand name: Voltaren®

### **Counterirritants**

Example brand names: IcyHot®, Biofreeze®

#### Common uses

- Provides temporary relief of minor aches and pain of muscles and joints
- Will numb the pain where the patch is placed.

#### **Dosing**

- Only use 1 patch at a time
- Clean and dry area before applying
- Apply for up to 8 hours
- Can apply new patch 3-4 times daily



#### Warning

- Do not use heating pad or warm gel pack with the patch
- Should not be used by children under 12 years old
- Seek medical attention if symptoms last longer than 7 days

#### Common uses

- Provides relief from pain associated with osteoarthritis in hand, wrist, elbow, foot, ankle, and knee
- Reduces inflammation and pain at application site

#### **Dosing**

- Apply to painful area following packaging directions
- Use for a maximum of 21 days

Note: It can take up to 7 days to experience relief after starting this gel

#### Warning

- Do not use on open wounds or with heating pads
- May increase the risk of heart and blood vessel problems
- Do not use if you have a history of asthma or allergic-type reactions with aspirin or other nonsteroidal medications (NSAID), such as ibuprofen and naproxen
- Using this product with oral NSAIDS could increase the risk of adverse effects

#### Common uses

- Provides temporary relief of aches and sprains of muscles and joints, back pain, and arthritis pain
- Relieves pain by stimulating the nerves with a warming and/or cooling feeling

#### Dosing

- Apply to painful area following packaging directions
- Use as needed for up to 7 days



#### Warning

- Use products only as directed
- Skin irritation or rash could occur
- Note that some products may contain lidocaine. Follow packaging directions.